

DMSO/Lidocaine 50%/0.5% Bladder Irrigation

Visually Impaired

If you are visually impaired and need help reading your label we can help. At your request, a prescription reader is available to patients in need. If you need help, please contact the pharmacy at 727-498-1228 and speak to a technician or pharmacist who will be able to advise you on your options and the cost. Additional charges do apply.

What is this medication?

This is a compounded bladder irrigation that contains Dimethyl Sulfoxide (DMSO) 50% and Lidocaine 0.5%. It is typically prescribed for the treatment of pain, swelling, and irritation of the bladder, or other forms of chronic bladder inflammation. DMSO reduces inflammation and pain, while Lidocaine provides local anesthetic relief to minimize discomfort during bladder instillation. If you have been given this drug for some other reason, talk with your doctor about the benefits and risks. Talk with your doctor if you have questions or concerns about using this drug.

What are the ingredients?

Dimethyl Sulfoxide (DMSO) 50% and Lidocaine 0.5%.

If you have any further questions or concerns about the ingredients used in your customized, compounded formulation, please contact one of our pharmacists at 727-498-1228.

What should I tell my healthcare provider before I take this medication?

If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. This drug may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

How should I use this medication?

This medication is instilled directly into the bladder by a healthcare professional, or at home if instructed by your provider. It is typically administered through a catheter and held in the bladder for 15–30 minutes, or as directed. Do not drink large volumes of fluids before installation. Follow all technique instructions carefully if administering at home.

What if I miss a dose?

If you miss an appointment or scheduled dose, contact your provider or pharmacist to reschedule. Do not attempt to self-administer without proper instruction. Do not double the dose to make up for a missed one.

What may interact with this medication?

Dimethyl Sulfoxide may decrease therapeutic effects of Dexrazoxane. *Avoid.* ([Lexi.com](#))
Dimethyl Sulfoxide may increase neurotoxic effects of Sulindac. Dimethyl Sulfoxide may decrease the metabolism of Sulindac. Specifically, the concentrations of the active sulfide metabolite are decreased. *Monitor.* ([Lexi.com](#)). Always check with your provider before starting or stopping any medication, including non-prescription products.

What side effect:

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Change in eyesight, eye pain, or severe eye irritation.

This list may not describe all possible side effects. Call your doctor for medical advice about side effects and any unusual problems while taking this medication. If you experience a serious side effect, you or your doctor may send a report to the Food and Drug Administration's (FDA) MedWatch Adverse Event Reporting program, online (www.fda.gov/Safety/MedWatch) or by phone (1-800-332-1088)

What should I watch while using this medication?

Notify your provider if symptoms worsen or new symptoms develop. Maintain good catheter hygiene if administering at home. Do not use it if the medication changes color or forms crystals. Report any signs of infection such as fever, foul-smelling urine, or discomfort.

Where should I keep my medication?

If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

Disposal: Do not flush it or pour it down the drain. The best option is to use a DEA-authorized drug take-back location, which you can find at rxdrugdropbox.org. If that's not available, mix the



medication (still in its vial or drawn into a syringe) with an undesirable substance like used coffee grounds or cat litter, seal it in a plastic bag, and discard it in the household trash—after removing any personal information from the label. Used syringes must be placed in an FDA-cleared sharps container and taken to an appropriate sharps disposal site. If you're unsure, contact your pharmacist at 727-498-1228 for guidance.

OFFER TO COUNSEL NOTICE

You are receiving this notice because you or your physician recently requested a compounded prescription to be filled by our pharmacy. You are entitled to receive medication counseling from one of our pharmacists regarding the following: Name and description of medication. Dose, route of administration, and duration of medication. Intended use of medication and expected action. Special directions and precautions for use. Common adverse effects or interactions. Techniques for self-monitoring. Proper storage. Prescription refill information.

You may contact a pharmacist at (727) 498-1228 Monday through Friday, 9AM to 5PM EST and Saturday 9AM-1PM EST.

PharmaLabs' notice of Privacy Practices may be viewed on the company website. You may view and acknowledge receipt of these offers or refuse your right to be counseled online at: <http://www.plabrx.com/offer>

Our pharmacists will be happy to discuss any additional questions or concerns you may have.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.