

## **Semaglutide/Cyanocobalamin (2.5mg/0.5mg)/mL Subcutaneous Injection**

### **Visually Impaired**

If you are visually impaired and need help reading your label we can help. At your request, a prescription reader is available to patients in need. If you need help, please contact the pharmacy at 727-498-1228 and speak to a technician or pharmacist who will be able to advise you on your options and the cost. Additional charges do apply.

### **What is this medication?**

This medication is a combination of Semaglutide, a GLP-1 receptor agonist, and Cyanocobalamin, a form of vitamin B12. It is used as part of a medically supervised weight loss program to help reduce appetite, support blood sugar control, and promote fat metabolism. Semaglutide helps regulate hunger and insulin release, while cyanocobalamin supports energy production and prevents B12 deficiency.

### **What are the ingredients?**

This is a compound formulation of Semaglutide 2.5 mg/mL and Cyanocobalamin 0.5 mg/mL (Vitamin B12).

If you have any further questions or concerns about the ingredients used in your customized, compounded formulation, please contact one of our pharmacists at 727-498-1228.

### **What should I tell my healthcare provider before I take this medication?**

Inform your provider if you have a history of thyroid cancer, pancreatitis, gallbladder disease, kidney problems, or diabetes. Also let them know if you are pregnant, planning to become pregnant, or breastfeeding. Tell your provider about any other medications or supplements you are taking.

### **How should I use this medication?**

Inject this medication subcutaneously (under the skin) once weekly or as directed by your healthcare provider. Preferred injection sites include the abdomen, thigh, or upper arm. Rotate injection sites each time to avoid irritation. Do not inject into muscle or vein. Follow instructions provided by your pharmacist or clinician on proper dose preparation and injection technique.

### **What if I miss a dose?**

If you miss a dose, take it as soon as you remember, unless it's close to the time of your next scheduled dose. In that case, skip the missed dose and resume your regular dosing schedule. Do not double up on doses. Contact your provider for guidance if you miss more than one dose.

### **What may interact with this medication?**

Semaglutide may slow digestion, which can affect the absorption of other oral medications. Tell your provider if you are taking oral birth control, blood sugar-lowering medications, or thyroid replacement therapy. Avoid alcohol, which may increase the risk of low blood sugar or gastrointestinal side effects. Always inform your healthcare provider before starting or stopping any medications or supplements.

### **What side effect:**

Common side effects may include nausea, vomiting, diarrhea, constipation, or decreased appetite. These usually lessen over time. Serious side effects may include pancreatitis, gallbladder issues, or allergic reactions. Cyanocobalamin is generally well tolerated, though mild redness at the injection site may occur.

This list may not describe all possible side effects. Call your doctor for medical advice about side effects and any unusual problems while taking this medication. If you experience a serious side effect, you or your doctor may send a report to the Food and Drug Administration's (FDA) MedWatch Adverse Event Reporting program, online ([www.fda.gov/Safety/MedWatch](http://www.fda.gov/Safety/MedWatch)) or by phone (1-800-332-1088)

### **What should I watch while using this medication?**

Monitor your weight, appetite, blood sugar levels (if diabetic), and any gastrointestinal symptoms. Report significant weight loss or persistent side effects to your healthcare provider. Stay hydrated and maintain a balanced diet as advised by your care team.

### **Where should I keep my medication?**

Store according to the label instructions. Refrigeration is typically required, unless otherwise noted. Do not freeze. This medication is a sterile injectable product and expires 28 days after the vial is first punctured, or the date on the prescription vial label, whichever date comes first. Keep away from light and heat. Keep out of reach of children and pet.

**Disposal:** Do not flush it or pour it down the drain. The best option is to use a DEA-authorized drug take-back location, which you can find at [rxdrugdropbox.org](http://rxdrugdropbox.org). If that's not available, mix the medication (still in its vial or drawn into a syringe) with an undesirable substance like used coffee grounds or cat litter, seal it in a plastic bag, and discard it in the household trash—after



removing any personal information from the label. Used syringes must be placed in an FDA-cleared sharps container and taken to an appropriate sharps disposal site. If you're unsure, contact your pharmacist at 727-498-1228 for guidance.

### **OFFER TO COUNSEL NOTICE**

You are receiving this notice because you or your physician recently requested a compounded prescription to be filled by our pharmacy. You are entitled to receive medication counseling from one of our pharmacists regarding the following: Name and description of medication. Dose, route of administration, and duration of medication. Intended use of medication and expected action. Special directions and precautions for use. Common adverse effects or interactions. Techniques for self-monitoring. Proper storage. Prescription refill information.

You may contact a pharmacist at (727) 498-1228 Monday through Friday, 9AM to 5PM EST and Saturday 9AM-1PM EST.

PharmaLabs' notice of Privacy Practices may be viewed on the company website. You may view and acknowledge receipt of these offers or refuse your right to be counseled online at: <http://www.plabrx.com/offer>

Our pharmacists will be happy to discuss any additional questions or concerns you may have.

**NOTE:** This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.