

## Kyzatrex 200mg Oral Capsule

### **Visually Impaired**

If you are visually impaired and need help reading your label we can help. At your request, a prescription reader is available to patients in need. If you need help, please contact the pharmacy at 727-498-1228 and speak to a technician or pharmacist who will be able to advise you on your options and the cost. Additional charges do apply.

### **What is this medication?**

Brand Name: Kyzatrex™

Generic Name: Testosterone undecanoate.

Kyzatrex™ is an oral testosterone replacement therapy (TRT) used in adult males with conditions associated with a deficiency or absence of endogenous testosterone (also known as hypogonadism). This includes: Primary hypogonadism (testicular failure) and Hypogonadotropic hypogonadism (pituitary or hypothalamic dysfunction).

### **What are the ingredients?**

Active Ingredient: Testosterone undecanoate 200 mg per capsule.

### **What should I tell my healthcare provider before I take this medication?**

Before using Kyzatrex™, tell your provider if you: Have prostate cancer or breast cancer. Have heart, liver, or kidney problems. Have sleep apnea. Have high red blood cell counts (polycythemia). Are taking insulin or oral diabetes medications. Have ever had a blood clot, heart attack, or stroke. Are taking anticoagulants or corticosteroids. Are planning to father a child — testosterone can reduce fertility.

### **How should I use this medication?**

Take exactly as prescribed, typically twice daily (morning and evening) with food.

Swallow capsules whole—do not chew or open them.

Must be taken with a meal that contains fat to ensure proper absorption.

Maintain consistent timing and dietary habits for each dose.

Do not exceed the prescribed dose.

### **What if I miss a dose?**

Take the missed dose as soon as you remember, if it's early in the day. If it's later in the day, skip the missed dose to avoid trouble sleeping. Do not double up on doses.

### **What may interact with this medication?**

Potential interactions include: Insulin or diabetes medications: may lower blood sugar. Blood thinners (e.g., warfarin): increased risk of bleeding. Corticosteroids: may increase risk of fluid retention. Other testosterone products or anabolic steroids.

Always check with your provider before starting or stopping any medication, including non-prescription products.

### **What side effect:**

**Common side effects:** Increased blood pressure. Headache. Acne. Increased red blood cell count. Enlarged prostate. Gastrointestinal symptoms (nausea, diarrhea)

**Serious side effects (seek immediate medical attention):** Chest pain or shortness of breath. Sudden vision or speech changes. Signs of liver problems (e.g., yellowing of skin or eyes). Swelling in legs, ankles, or feet. Changes in urination or prostate symptoms. Mood changes or aggressive behavior

This list may not describe all possible side effects. Call your doctor for medical advice about side effects and any unusual problems while taking this medication. If you experience a serious side effect, you or your doctor may send a report to the Food and Drug Administration's (FDA) MedWatch Adverse Event Reporting program, online ([www.fda.gov/Safety/MedWatch](http://www.fda.gov/Safety/MedWatch)) or by phone (1-800-332-1088)

### **What should I watch while using this medication?**

Monitor blood pressure regularly.

Routine blood tests may be required to check testosterone levels, liver function, cholesterol, hematocrit, and prostate-specific antigen (PSA).

Inform your doctor if you develop trouble breathing during sleep (possible sleep apnea).

Periodically evaluate for prostate health, especially if over age 40.

### **Where should I keep my medication?**

Store at room temperature (20°C to 25°C / 68°F to 77°F). Keep the bottle tightly closed and away from moisture or excessive heat. Keep out of reach of children and pets.

**Caution:** ***"Federal law prohibits the transfer of this drug to any person other than the patient for whom it was prescribed."***

**Disposal:** Do not flush or pour unused medication down the drain. The best option is to use a DEA-authorized drug take-back location, which you can find at [rxdrugdropbox.org](http://rxdrugdropbox.org). If that's not available, mix the medication (still in its vial or drawn into a syringe) with an undesirable substance like used coffee grounds or cat litter, seal it in a plastic bag, and discard it in the household trash—after removing any personal information from the label. Used syringes must be placed in an FDA-cleared sharps container and taken to an appropriate sharps disposal site. If you're unsure, contact your pharmacist at 727-498-1228 for guidance.

## **OFFER TO COUNSEL NOTICE**

You are receiving this notice because you or your physician recently requested a compounded prescription to be filled by our pharmacy. You are entitled to receive medication counseling from one of our pharmacists regarding the following: Name and description of medication. Dose, route of administration, and duration of medication. Intended use of medication and expected action. Special directions and precautions for use. Common adverse effects or interactions. Techniques for self-monitoring. Proper storage. Prescription refill information.

You may contact a pharmacist at (727) 498-1228 Monday through Friday, 9AM to 5PM EST and Saturday 9AM-1PM EST.

PharmaLabs' notice of Privacy Practices may be viewed on the company website. You may view and acknowledge receipt of these offers or refuse your right to be counseled online at: <http://www.plabrx.com/offer>

Our pharmacists will be happy to discuss any additional questions or concerns you may have.

**NOTE:** This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.